

Differential expression of laminin isoform ($\alpha 2$), integrins ($\alpha 3\beta 1$ and $\alpha 6\beta 4$) and cytokeratin 20 in *H. pylori* gastritis

M.G.F. Sales¹, L.E. Nasciutti¹, D.E. Lorena¹, M. Muzzi⁴ and L.C. Porto^{3,4}

¹Department of Histology and Embryology, UFRJ, Rio de Janeiro, ²School of Medicine Souza Marques, Rio de Janeiro, ³Department of Histology and Embryology, UERJ, Rio de Janeiro and ⁴Labs. Patologia Clínica e Investigação Ltda, Rio de Janeiro, Brazil

Summary. The expression of laminin-1 chains ($\beta 1$ and $\gamma 1$), laminin-2 (merosin), integrin receptors to laminin ($\alpha 3\beta 1$ and $\alpha 6\beta 4$) and cytokeratin (CK20) were studied by immunohistochemical methods in gastric biopsies from antrum of 25 patients. *H. pylori* gastritis was found in 19 cases and intestinal metaplasia (IM) in four from these 19. Another 13 biopsies, all with IM were immunostained to laminin-2. Laminin-1 chains in normal and gastritis areas without IM were expressed as a strong, linear and continuous deposit in the basement membranes of the superficial and glandular epithelium. In metaplastic glands the reactivity to laminin-1 chains was decreased. Merosin was discontinuous when a moderate to accentuated *H. pylori* glandular colonization was present. Samples with IM were negative to laminin-2. The $\alpha 3\beta 1$ and $\alpha 6\beta 4$ integrins were negative only in IM gastric biopsies. The CK20 immunoreactivity was strong and homogeneous in the cells at the tip and the upper portion of foveolae in normal areas and in gastritis with IM the reactivity to CK 20 was heterogeneous. A differential expression of laminin isoforms is related to inflammation and subsequent IM caused by *H. pylori*. The alterations of $\alpha 3\beta 1$ and $\alpha 6\beta 4$ parallel both modifications in merosin and CK20 expression in *H. pylori* chronic gastritis.

Key words: Laminins chains, Cytokeratin, *Helicobacter pylori* gastritis, Integrins

Introduction

The processes of development, cell growth and cell differentiation are closely related with the interactions between cells and the extracellular matrix, which is composed by interstitial components and basement membrane (BM) (Ekblom et al., 1986). BMs are

specialized, sheet-like, extracellular matrix structures that separate the connective tissue from epithelia, muscle fibers, blood vessels and nerves (Foidart et al., 1980; Lissitzky et al., 1986; Edds, 1997). In recent years, the importance of extracellular matrix components, especially those located at the epithelial-mesenchymal interface, has been illustrated in the developing human small intestine (Haffen et al., 1989; Beaulieu et al., 1991). Laminins are the major BM glycoproteins throughout the body, composed by 3 chains ($\alpha 1$ -5; $\beta 1$ -3; $\gamma 1$ -2) assembled into heterotrimeric isoforms (Chung et al., 1979; Timpl et al., 1979; Burgeson et al., 1994; Miner et al., 1997). Laminins arise very early during the development, and play an important role in the interactions and in the maintenance of gland and epithelia adult state (Beaulieu and Vachon, 1994; Simon-Assmann et al., 1994; Perreault et al., 1995). In particular, merosin, or laminin $\alpha 2$ chain, is restricted to developing the pit-gland structure of the forming glands during gastric development up 20 weeks of gestation (Tremblay and Ménard, 1996), and in the normal adult state is confined to the gastric and intestinal glandular basement membranes (Simo et al., 1991; Virtanen et al., 1995; Tremblay and Ménard, 1996). However, little is known about the composition and distribution of the different BM components associated with the developing gastric mucosa and mature gastric glands (Tremblay and Ménard, 1995, 1996).

Integrins are heterodimeric cell surface glycoproteins that mediate divalent cation-dependent cell-cell and cell-matrix interactions (Buck and Horwitz, 1987; Hynes, 1987). They are composed of a single α subunit associated with a single β subunit and this association is required for cell surface expression. The interactions mediated by integrins are responsible for certain typical properties of adhesive cells, such as attachment and migration, but these molecules are also recognized to contribute to intracellular signaling processes (Hynes, 1992; Humphries, 1999). Amongst the laminin binding integrins, the $\alpha 3\beta 1$ integrin binds laminin 2, 4 and 5 (Rousselle and Aumailley, 1994) and cannot play a major role in cell adhesion to laminin-1 (Aumailley et al., 1990). The integrin $\alpha 6\beta 4$ has been

Offprint requests to: Prof. Luís Cristóvão Porto, Department of Histology and Embryology, IBRAG, UERJ, Av. Prof. Manoel de Abreu, 444 30 andar, 20551-170 Rio de Janeiro, Brazil. Fax: 55 21 5876511. e-mail: lcporto@uerj.br

studied by immunoelectron microscopy and its distribution has been described at the basal side of several epithelial cell types, suggesting that this integrin plays a role in polarity and adhesion of epithelial cells to the BM (Marchisio et al., 1991). Several works demonstrated that $\alpha 6 \beta 4$ polarization is lost upon tumorigenic transformation of epithelial cells (Rabinovitz and Mercurio, 1996).

Keratin is the most complex of intermediate filaments and belongs to a great family of related proteins that vary in molecular weight and are coded by a complex set of genes, expressed in different types of epithelium that contain distinct keratin profiles (Franke et al., 1981). Cytokeratin 20 (CK20), was specifically detected as a component of intestinal and gastric foveolar epithelium (Moll et al., 1990, 1993) and remains unchanged even during intestinal metaplasia or malignant progression (Moll et al., 1993; Schwerer and Baczako, 1996).

H. pylori has been considered as a causal agent of chronic gastritis in humans and may be associated with gastric cancer (Valkonen et al., 1994; Solcia et al., 1996). Several works have emphasized that the adhesion of *H. pylori* to laminin disorganizes the gastric epithelium altering the interactions between laminin and laminin receptor in gastric epithelial cells (Slomiany et al., 1991; Valkonen et al., 1994; Sales et al., 1998). The mucosal BM has been considered to play a role in the high capacity of mucosal repair after injury (Mikami et al., 1994), but the way gastric BM component expression occurs during injury, remains unknown.

We decided to detect, by immunohistochemical methods, the expression of $\beta 1$, $\gamma 1$, and $\alpha 2$ laminin chains, their integrin receptors $\alpha 6 \beta 4$ and $\alpha 3 \beta 1$ and cytokeratin 20 (CK20) in the gastric mucosa with *H. pylori* chronic gastritis. Aim: to investigate whether the inflammatory and metaplastic process caused by *H. pylori* alters the expression of BM extracellular molecules and CK20 intermediate filament.

Materials and methods

Gastric samples

Samples of antral gastric mucosa from 25 patients with chronic gastritis were obtained by endoscopic means at the University Hospital of UFRJ, Rio de Janeiro. Patients ranged from 37 to 67 years old, being 18 male and 9 female. The principal symptoms reported were: abdominal pain, epigastric burning and distension, periodic nausea, flatulence and halitosis. The criteria of selection were: a) *H. pylori* negative cases were included based on no previous eradication therapy; b) patients with neoplasia, especially in the gastrointestinal tract, were excluded; c) all the patients had not a specific drug intake, especially non-steroid anti-inflammatory drugs, at least 2 weeks before the endoscopy; d) cases with previous gastric surgery were also excluded. Control fragments were obtained from four patients with focal

enanthematos areas in which the histopathology revealed only minimal gastritis; the samples collected from normal endoscopic adjacent areas exhibited normal histology. From each sample area ($n = 29$), one fragment was immediately frozen in liquid nitrogen and stored at -70°C , the other being embedded in paraffin. Diagnoses were confirmed from standard hematoxylin-eosin, Alcian-Blue-PAS and Waysson method for *H. pylori*. Another 13 antral biopsies in which chronic gastritis with intestinal metaplasia without dysplasia was histologically detected, were randomly selected from paraffin-embedded samples of the Labs Patologia Clínica e Investigação Laboratorial pathology service. Gastritis was evaluated based on the criteria of the Sydney score (Sipponen et al., 1991).

Antibodies

The monoclonal antibodies used in this study were against laminin chains: $\alpha 2$ (Mab1922 diluted 1:1000), $\beta 1$ (Mab1928 diluted 1:500), $\gamma 1$ (Mab1920 diluted 1:500), from Chemicon International Inc, against integrin receptor to laminins: $\alpha 6 \beta 4$ (Mab1982 diluted 1:100) from Chemicon International Inc. and $\alpha 3 \beta 1$ (MO608 diluted 1:200) from Dako Corporation. All antibodies were used for immunofluorescence staining on frozen sections and meroisin ($\alpha 2$ laminin chain) was also applied in paraffin sections. For the cytoskeleton, the monoclonal antibody anti-cytokeratin 20 (CK 20) from Dako Corporation was used on the paraffin sections.

Immunohistochemistry

The frozen tissue specimens from normal and chronic gastritis areas, were sectioned at $5\ \mu\text{m}$ and fixed in acetone precooled to -20°C . After washing in phosphate buffer saline/serum bovine albumin (PBS/BSA), pH 7.4, the frozen sections were incubated with the primary antibody (anti- $\alpha 2$, $\beta 1$, $\gamma 1$ laminin chains, anti- $\alpha 3 \beta 1$ or $\alpha 6 \beta 4$ integrin) for one hour. The procedure was continued by incubating the sections with a fluorescein-isothiocyanate-conjugated goat anti-mouse, for 30 minutes, followed by three washes in PBS and mounting in buffered glycerol. The specimens were examined in a Zeiss microscope with epi-illumination and the appropriate filter.

For the immunoperoxidase staining, the paraffin-embedded specimens were sectioned at $4\ \mu\text{m}$ on silanized slides to avoid damage to the sections during the procedure. A duet kit (Dako Corporation) was used according to the protocols provided by the distributor. To detect CK20, the sections were dewaxed in xylene and rehydrated with graded alcohol to distilled water. Antigen retrieval was carried out in conjunction with microwave oven heating. Following washing for 10 minutes in PBS at pH 7.4, the endogenous peroxidase activity was blocked with 0.6% H_2O_2 in methanol for 15 minutes. The sections were then washed in PBS for 10

minutes and incubated with non-immune sheep serum at 1% to reduce the background. After this, the primary antibody was applied (anti- $\alpha 2$ laminin chain or anti-CK20), diluted with non-immune sheep-serum at 1%, and incubated for one hour. For the evaluation of immunostaining, the primary antibody was replaced by non-immune serum to provide a negative control. This was followed by three washes in PBS and a second incubation with biotinylated goat anti-mouse antibody for 30 minutes. Following this, the sections were incubated with streptavidin-peroxidase complex for 30 minutes. All the incubations were carried out at room temperature. The immunoreactivity was detected by a

substrate solution containing H_2O_2 and 3,3' diaminobenzidine (DAB) in PBS, pH 7.4. The sections were then counterstained with Mayer's hematoxylin (Merck) diluted with distilled water 1:2, washed in water, dehydrated in alcohol, cleared in xylene and mounted with Entellan (Merck).

Results

From the 25 samples of chronic antral gastritis, 19 (76%) were positive to *H. pylori*, having quantities that ranged from minimal (36%) to moderate (36%) and accentuated (4%). In addition, from the total number of




Fig. 1. Immunofluorescence to laminin-1 chains of normal gastric mucosa areas and of *H. pylori* chronic gastritis with intestinal metaplasia. **A and B.** FITC indirect immunofluorescence with anti- $\beta 1$ chain antibody. **C and D.** FITC indirect immunofluorescence with anti- $\gamma 1$ chain antibody. **A.** Superficial epithelium (*) basement membrane from normal antrum area showing an intense, linear and continuous reaction to $\beta 1$ chain (arrow-heads). Basal lamina of capillaries (thin arrows) in the stroma also exhibit intense staining. **B.** Chronic gastritis with intestinal metaplasia. Weak to moderate and irregular reaction to $\beta 1$ chain in the basement membrane (arrow-heads) underneath the glandular portion with three goblet cells (*). Basal lamina of a capillar (thin arrow) in the stroma exhibit intense staining. **C.** Superficial epithelium (*) basement membrane from normal antrum area showing an intense and continuous reaction to $\gamma 1$ chain (arrow-heads). **D.** Chronic gastritis with intestinal metaplasia. Irregular and discontinuous reaction to $\gamma 1$ chain in the basement membrane (arrow-heads) of a intestinalized gland portion. x 860

samples, 4 showed intestinal metaplasia: 2 complete, exhibiting all the cellular components of the intestinal mucosa, and 2 incomplete.

Expression of $\beta 1$ and $\gamma 1$ laminin chains

Laminin $\beta 1$ chain was observed on the BM of superficial (Fig. 1A) and glandular epithelium, from normal areas, as a linear and continuous deposit. Those areas, histologically diagnosed as gastritis, showed that, independently of whether or not *H. pylori* was detected, and independently of gastritis activity and the extent of inflammation, the BMs were also stained with a linear and continuous deposit of $\beta 1$ chain (not shown). The expression of laminin $\gamma 1$ was similar to $\beta 1$ chain showing the same pattern of staining (Fig. 1C). However, in the four cases of gastritis with metaplasia, where the inflammation ranged from moderate to

accentuated, the reactivity was less intense and staining discontinuity of the BMs was seen both with anti- $\beta 1$ (Fig. 1B) and $\gamma 1$ (Fig. 1D) antibodies in these metaplastic cases. Reactivity to basal lamina of capillaries was intense with both antibodies.

Expression of $\alpha 2$ laminin chain (merosin)

In the normal areas, the BMs of the glandular epithelium showed a linear and continuous deposit of $\alpha 2$ chain (Fig. 2A). However, in the gastritis areas, the expression of $\alpha 2$ chain was not uniform. The glandular epithelium BMs showed staining patterns varying from linear continuous to discontinuous. In the samples with gastritis where no *H. pylori* was detected, the BMs showed a linear and continuous deposit of $\alpha 2$ chain and discontinuity was seen in those samples exhibiting moderate and accentuated presence of *H. pylori* and

Table 1. Immunoreactivity of pyloric glands to basement membrane laminin isoforms, to cell membrane integrin receptors, and of superficial epithelium to cytokeratin 20 (CK20) in gastric fragments of patients with chronic gastritis.

	LAMININS $\beta 1$ AND $\gamma 1$	LAMININ $\alpha 2$	INTEGRINS $\alpha 3\beta 1$ and $\alpha 6\beta 4$	CK20 SUPERFICIAL EPITHELIUM
Normal antrum and chronic gastritis without <i>H. pylori</i> (no inflammatory activity)	Intense, continuous and linear	Intense, continuous and linear	Intense baso lateral deposit. Moderate in chronic gastritis	Intense baso-apical cytoplasmatic filaments
Chronic gastritis with moderate or accentuated <i>H. pylori</i> colonization (moderate and accentuated inflammatory activity)	Intense, continuous and linear	Intense, discontinuous and linear	$\alpha 3\beta 1$ – moderate to intense basolateral deposit $\alpha 6\beta 4$ – irregular some glands with basal others with lateral deposits	Intense basal cytoplasmatic filaments
Intestinal metaplasia areas of <i>H. pylori</i> chronic gastritis (moderate and accentuated inflammatory activity)	Weak to moderate, continuous and linear	Non-reactive	Non-reactive	Goblet cells – non-reactive Absorptive columnar cell – apical pole




Fig. 2. Immunostaining to laminin $\alpha 2$ chain of *H. pylori* chronic gastritis with or without intestinal metaplasia. **A.** FITC indirect immunofluorescence on frozen section. Chronic gastritis without intestinal metaplasia. The glandular basement membranes show an intense, linear and continuous deposit. x 215. **B.** Immunoperoxidase method on paraffin embedded section. No reaction is seen in the chronic gastritis with intestinal metaplasia. x 860

inflammatory activity. No reaction was seen in the four cases of intestinalized glands, but in all the 25 samples, strong stromal reactivity was found.

The sections of paraffin-embedded samples were also stained by peroxidase method with anti-merosin antibody. The results were similar to those with the immunofluorescence method (not shown). No reactivity of gastric BM areas with intestinal metaplasia to merosin was confirmed on a further 13 paraffin-embedded samples. The intestinalized glands from complete or incomplete forms were always negative to merosin. In contrast, the normal glands had a linear reaction to $\alpha 2$ (Fig. 2B).

Expression of cytokeratin 20

Paraffin-embedded sections were stained to CK20, in order to ascertain whether any alteration would be seen in the cytoskeleton. Immunoreactivity was restricted to the superficial epithelium, upper foveolae and to a variable number of single CK20-positive epithelial cells in the upper zone of the normal pyloric glands (Fig. 3A). In normal areas, the strong reactivity was basal, with a filamentous pattern with a predominantly baso-apical orientation. Near the lateral and basal membranes, a reinforcement of immunoreactivity was observed (Fig. 3B). The same staining was seen in the cases with gastritis without metaplasia. Intestinal metaplasia areas exhibited a different pattern of staining. The reactivity was irregular and heterogeneous, i.e., some cells showed an intense reaction while others were negative. Goblet cells did not react with the CK20 antibody and or absorptive columnar cells, the reactivity was restricted to the apical cell pole and basal reinforcement was seen, as in the samples without metaplasia (Fig. 3C).

Expression of $\alpha 3\beta 1$ and $\alpha 6\beta 4$ integrins receptors to laminin

In order to investigate the presence of integrin receptors to laminin in chronic gastritis, antibodies against $\alpha 3\beta 1$ and $\alpha 6\beta 4$ were utilized in cryosections stained by indirect immunofluorescence. The $\alpha 3\beta 1$ integrin was detected in all the normal gastric samples as an intense basolateral deposit of the pyloric glands and also in endothelial and interstitial cells (Fig. 4A). In the sections with chronic gastritis and *H. pylori* colonization, the reaction ranged from negative in the metaplastic glands to a moderate to strong reactivity in non-intestinalized epithelium, independently of the inflammatory activity. The absence of reactivity in intestinalized glands was not associated with the type of




Fig. 3. Immunoperoxidase to CK20 on paraffin embedded section. **A.** Pyloric glands showing few positive CK20 cells (arrows). **B.** Superficial epithelium of normal area with a regular distribution of CK20. **C.** Chronic gastritis with intestinal metaplasia exhibiting an irregular aspect of CK20 distribution. Goblet cell did not react with CK20 antibody (*). $\times 860$




Fig. 4. Immunofluorescence to integrins receptors to laminin of normal gastric mucosa areas and of *H. pylori* chronic gastritis with and without intestinal metaplasia. **A.** $\alpha3\beta1$ integrin in glandular epithelium from normal area showing a strong basolateral reaction (arrows). Endothelial and some interstitial cells in the stroma exhibit intense staining. **B.** $\alpha6\beta4$ in chronic gastritis. Moderate reaction is seen in the basolateral portion of the glandular epithelium (arrow-heads). **C.** $\alpha6\beta4$ in chronic gastritis with intestinal metaplasia. No staining in the glandular basement membrane is observed. $\times 860$

intestinal metaplasia. The expression of $\alpha6\beta4$, although similar to that seen in $\alpha3\beta1$ staining, in some cases showed a lateral reaction and while in others, a basal localization (Fig. 4B). Areas with intestinal metaplasia did not react with the $\alpha6\beta4$ antibody (Fig. 4C).

The results were summarized on Table 1.

Discussion

In this report, we analyzed the expression and distribution of $\alpha2$, $\beta1$ and $\gamma1$ laminin chains, their $\alpha3\beta1$ and $\alpha6\beta4$ integrin receptors and the cytokeratin 20 of antral biopsies with gastritis with or without *H. pylori* cells, and correlated their distribution with morphological changes during this disease.

The patterns of $\beta1$ and $\gamma1$ expression in normal areas are similar to those reported for the human fetal and adult gastric mucosa (Virtanen et al., 1995; Tremblay and Ménard, 1996). Since the basement membranes were also stained with a linear and continuous deposit of $\beta1$ and $\gamma1$ laminin chains, gastritis with *H. pylori* and subsequent inflammation not change the expression of these two laminin chains. However, the intestinalization process and the degree of inflammation suggest a modification of the expression of these two laminin chains. In contrast, in areas from gastritis, the merosin reactivity ranged from continuous to discontinuous, and the discontinuity was always *H. pylori* associated, thus indicating that $\alpha2$ chain expression can be modulated by *H. pylori* colonization and by the amount of inflammatory cells present in the lamina propria. The results of this study also showed that when gastric glands lose their typical appearance and acquire an intestinal pattern, the expression of merosin is removed from the BMs. Moreover, even in absence of merosin in the BM intestinalized glands, the stroma stained to it. Then, in gastritis with complete or incomplete intestinal metaplasia, merosin expression may not be regulated by heterologous cell-cell interaction as suggested by Simon-Assman et al. (1994). The interactions with epithelial and mesenchyma-derived cells, on producing merosin, may be dependent on the maintenance of the differentiated state of both cellular types (Parker et al., 1974; Neal and Potten, 1981; Keding, 1994). Merosin is not only involved with glandular development, but also with the maintenance of adult glandular structures of all gastrointestinal tract (Simo et al., 1991; Simon-Assman et al., 1994; Perrault et al., 1995; Virtanen et al., 1995; Lohi et al., 1996; Tremblay and Ménard, 1996).

Although intestinal metaplasia develops after *H. pylori* infection, it may also appear for other reasons. The fact that a mucosal colonization by *H. pylori* may change the merosin expression (discontinuity), suggests that the binding of *H. pylori* to surface epithelial cells and subsequent inflammatory response play a role in the modulation of this molecule during the process of glandular renewal.

Intestinal metaplasia is a different tissue to the non-metaplastic gastric glands and expresses different antigens to non-metaplastic epithelium. Several works have suggested that the development of gastric adenocarcinoma is related to intestinal metaplasia (Lauren, 1991). Furthermore, intestinal-type dysplasias, adenomas and also adenocarcinomas arise from this region, suggesting that they share a common origin (Hattori and Sugihara, 1996). It seems to us that in the absence of merosin during the intestinalization process, the stem cell loses its pathways of differentiation and this loss may leave the glandular cell in a determined tubule to a preneoplastic state.

The integrins $\alpha 6 \beta 4$ and $\alpha 3 \beta 1$ have been related to an anchorage process between the extracellular matrix and the actin or keratin-based cytoskeleton (Carter et al., 1991; Niessen et al., 1997). The $\alpha 6 \beta 4$ integrin mediates stable anchorage of cells to laminins (Niessen et al., 1997) while the $\alpha 3 \beta 1$ integrin plays a role in matrix assembly (DiPersio et al., 1997). In addition, studies relate the loss of polarization (Tennenbaum et al., 1993) or its absence (Downer et al., 1993; Gui et al., 1995) in the tumoral process. In this work $\alpha 6 \beta 4$ was detected at basal localization and also at the lateral membranes in glands colonized by *H. pylori*. In intestinalized glands, no reaction to $\alpha 6 \beta 4$ and $\alpha 3 \beta 1$ integrins was observed, as in the merosin staining. Previously described as a laminin-1 receptor (Lee et al., 1992) it has now been confirmed that $\alpha 6 \beta 4$ is a receptor not only for laminin-1, but also for several laminin isoforms (Mercurio, 1995; Spinard et al., 1995). No reactivity of either merosin or $\alpha 6 \beta 4$ in the glands displaying intestinal metaplasia, suggests that this integrin is also a merosin receptor. The results of this work are partially different that those found by Tani et al. (1996), in which $\alpha 6 \beta 4$ was detected in glands with intestinal metaplasia, although like this study, no merosin was observed in intestinalized glands neither in the complete or incomplete forms.

The results of CK20 showed that in normal areas and in gastritis without metaplasia the immunoreactivity was strong and homogeneous in the cells at the tip and the upper portion of the foveolae. In the lamina propria, pyloric glands, few cells positive to CK20 were exhibited that were probably endocrine cells, as reported by Moll et al. (1993). However, in gastritis with intestinal metaplasia, the reactivity to CK 20 was not homogeneous, showing some cells with intense immunolabeling while others were negative. This difference can be explained by the mosaic-like patterns of positive and negative cells, which is a feature of the distribution of CK20 in normal and transformed tissues

(Moll et al., 1990). Schwerer and Baczako (1996) reported an intense and uniform reactivity to CK20 in enterocytes and goblet cells in the same disease. We did not find a positive reaction in goblet cells, probably by the density of intermediate filaments is low in central portions of these cells, due to mucin accumulation, as reported by Specian and Neutra (1984). Moreover, the pattern of expression of CK20 in association with CK7 was recently used to distinguish Barret's esophagus from intestinal metaplasia (Ormsby et al., 1999). In the intestinal metaplasia, cells positive for CK20, alternating with negative cells, probably differ functionally from each other.

Taken together, these findings suggest that a differential expression of laminin isoforms is related to inflammation and subsequent intestinal metaplasia caused by *H. pylori*. In conclusion, the alterations of integrins ($\alpha 3 \beta 1$ and $\alpha 6 \beta 4$) parallel both modifications in laminin isoform ($\alpha 2$) and CK20 expression in *H. pylori* chronic gastritis.

References

- Aumailley M., Timpl R. and Sonnenberg A. (1990). Antibody to integrin $\alpha 6$ subunit specifically inhibits cell-binding to laminin fragment 8. *Exp. Cell Res.* 188, 55-60.
- Beaulieu J.-F. and Vachon P.H. (1994). Reciprocal expression of laminin A-chain isoforms along the crypt-villus axis in the human small intestine. *Gastroenterology* 106, 829-839.
- Beaulieu J.-F., Vachon P.H. and Chartrand S. (1991). Immunolocalization of extracellular matrix components during organogenesis of the human small intestine. *Anat. Embryol.* 183, 363-369.
- Buck C. and Horwitz A.F. (1987). Cell surface receptors for extracellular matrix molecules. *Annu. Rev. Cell Biol.* 3, 179-205.
- Burgeson R.E., Chiquet M., Deutzmann R., Ekblom P., Engel J., Kleiman H., Martin G.R., Meneguzzi G., Paulsson M., Sanes J., Timpl R., Tryggvason K., Yamada Y. and Yurchenco P.D. (1994). A new nomenclature for the laminins. *Matrix Biol.* 14, 209-211.
- Carter W.G., Ryan M.C. and Gahr P.J. (1991). Epiligrin, a new cell adhesion ligand for integrin $\alpha 3 \beta 1$ in epithelial basement membranes. *Cell* 65, 599-610.
- Chung A.E., Jaffe R., Freeman I.L., Vergnes J.P., Braginski J.E. and Carlin B. (1979). Properties of a basement membrane-related glycoprotein synthesized in culture by a mouse embryonal carcinoma-derived cell line. *Cell* 16, 277-287.
- DiPersio C.M., Hodivala-Dilke K.M., Jaenisch R., Kreidberg J.A. and Hynes R.O. (1997). $\alpha 3 \beta 1$ integrin is required for normal development of the epidermal basement membrane. *J. Cell Biol.* 137, 729-742.
- Downer C.S., Watt F.M. and Speight P.M. (1993). Loss of $\alpha 6$ and $\beta 4$ integrin subunits coincides with the loss of basement membrane components in oral squamous cell carcinomas. *J. Pathol.* 171, 183-190.
- Edds L.L. (1997). Laminin and the mechanism of neuronal outgrowth. *Brain Res. Rev.* 23, 1-27.
- Ekblom P., Vestweber D. and Kemler R. (1986). Cell matrix interactions and cell adhesion during development. *Annu. Rev. Cell Biol.* 2, 27-47.
- Foidart J.-M., Bere E.W., Yaar Jr M., Rennard S.I., Gullino M., Martin G.R. and Katz S.I. (1980). Distribution and immunoelectron

Laminins and CK20 in *H. pylori* gastritis

- microscopic localization of laminin, a noncollagenous basement membrane glycoprotein. *Lab. Invest.* 42, 336-342.
- Franke W.W., Schiller D.L., Moll R., Winter S., Schmid E., Engelbrecht I., Denk H., Krepler R. and Platzer B. (1981). Diversity of cytokeratins: differentiation-specific expression of cytokeratin polypeptides in epithelial cells and tissues. *J. Mol. Biol.* 153, 933-959.
- Gui G.P., Puddefoot J.R., Vinson G.P., Wells C.A. and Carpenter R. (1995). Modulation of very late activation-2 laminin receptor function in breast cancer metastasis. *Surgery* 118, 245-250.
- Haffen K., Keding M. and Simon-Assman P. (1989). Cell contact dependent regulation of enterocytic differentiation. In: Human gastrointestinal development. Leberthal E. (Ed). Raven press. New York. pp 19-39.
- Hattori T. and Sugihara H. (1996). The pathological sequence in the development of gastric cancer: I. *Scand. J. Gastroenterol.* 31, 34-35.
- Humphries M.J. (1999). Towards a structural model of an integrin. *Biochem. Soc. Symp.* 65, 63-78.
- Hynes R.O. (1987). Integrins: a family of cell surface receptors. *Cell* 48, 549-554.
- Hynes R.O. (1992). Integrins: versatility, modulation, and signaling in cell adhesion. *Cell* 69, 11-25.
- Keding M. (1994). Differential expression of laminin isoforms in the developing human and mouse intestine. *Dev. Dyn.* 201, 71-85.
- Lauren P. (1991). Histogenesis of intestinal and diffuse types of gastric carcinoma. *Scand. J. Gastroenterol.* 180, 160-164.
- Lee E.C., Lotz M.M., Steele Jr. G.D. and Mercurio A.M. (1992). The integrin alpha 6 beta 4 is a laminin receptor. *J. Cell Biol.* 117, 671-678.
- Lissitzky J.C., Kopp F., Charpi C., Chiarotti Y., Bignon C. and Martin P.M. (1986). Laminine: Biosynthèse, structure et fonctions. *Path. Biol.* 34, 995-963.
- Lohi J., Leivo I., Tani T., Kiviluoto T., Kivilaakso E., Burgeson R.E. and Virtanen I. (1996). Laminins, tenascin and type VII collagen in colorectal mucosa. *Histochem. J.* 28, 431-440.
- Marchisio P.C., Bondanza S., Cremona O., Cancedda R. and De Luca M. (1991). Polarized expression of integrin receptors ($\alpha 6 \beta 4$, $\alpha 2 \beta 1$, $\alpha 3 \beta 1$ and $\alpha v \beta 5$) and their relationship with the cytoskeleton and basement membrane matrix in cultured human keratinocytes. *J. Cell Biol.* 112, 761-763.
- Mercurio A.M. (1995). Laminin receptors: achieving specifically through cooperation. *Trends Cell Biol.* 5, 419-423.
- Mikami H., Watanabe S., Hirose M. and Sato N. (1994). Role of extracellular matrix in wound repair by cultured gastric mucosal cells. *Biochem. Biophys. Res. Commun.* 202, 285-292.
- Miner J.H., Patton B.L., Lentz S.L., Gilbert D.J., Snider W.D., Jenkins N.A., Copeland N.G. and Sanes J.R. (1997). The laminin α chains: expression, developmental transitions, and chromosomal locations of $\alpha 1$ -5, identification of heterotrimeric laminins 8-11, and cloning of a novel $\alpha 3$ isoform. *J. Cell Biol.* 137, 685-701.
- Moll R., Schiller L. and Franke W.W. (1990). Identification of protein IT of the intestinal cytoskeleton as a novel type I cytokeratin with unusual properties and expression patterns. *J. Cell Biol.* 111, 567-580.
- Moll R., Zimbelmann R., Goldschmidt D., Keith M., Laufer J., Kasper M., Koch P.J. and Franke W.W. (1993). The human gene encoding cytokeratin 20 and its expression during fetal development and in gastrointestinal carcinomas. *Differentiation* 53, 75-93.
- Neal J.V. and Potten C.S. (1981). Circadian rhythms in the epithelial and the pericryptal fibroblast sheath in three different sites in the murine intestinal tract. *Cell Tissue Kinet.* 14, 581-587.
- Niessen C.M., Huisman E.H., Rots E.S., Sanchez-Aparicio P. and Sonnenberg A. (1997). Integrin $\alpha 6 \beta 4$ forms a complex with the cytoskeletal protein HD1 and induces its redistribution in transfected COS-7 cells. *Mol. Biol. Cell* 8, 555-566.
- Ormsby A.H., Goldblum J.R., Rice T.W., Richter J.E., Falk G.W., Vaezi M.F. and Gramlich T.L. (1999). Cytokeratin subsets can reliably distinguish Barrett's esophagus from intestinal metaplasia of the stomach. *Hum. Pathol.* 30, 288-294.
- Parker F.G., Barnes E.N. and Kaye G.I. (1974). The pericryptal fibroblast sheath. IV. Replication, migration and differentiation of the subepithelial fibroblasts of the crypt and villus of the rabbit jejunum. *Gastroenterology* 67, 607-621.
- Perreault N., Vachon P. H. and Beaulieu J-F. (1995). Appearance and distribution of laminin A chain isoforms and integrin $\alpha 2$, $\alpha 3$, $\alpha 6$, $\beta 1$, and $\beta 4$ subunits in the developing human small intestinal mucosa. *Anat. Rec.* 242, 242-250.
- Rabinovitz I. and Mercurio A.M. (1996). The integrin $\alpha 6 \beta 4$ and the biology of carcinoma. *Biochem. Cell Biol.* 74, 811-821.
- Rousselle P. and Aumailley M. (1994). Laminin is more efficient than laminin in promoting adhesion of primary keratinocytes and some other epithelial cells and has a different requirement for integrin receptors. *J. Cell Biol.* 125, 205-214.
- Sales M.G.F., Lellis J.C.B., Sobral M.F.P.S., Muzzi M., Lorena D.E., Ferreira F.C.C., Nasciutti L.E., Vargas C. and Porto L.C.M. (1998). Laminin, $\beta 1$, $\alpha 1$ and $\alpha 2$ laminin chains and integrin $\alpha 6 \beta 4$ distributions in *Helicobacter pylori* gastritis. *Digestion* 59, 390.
- Schwerer M.J. and Baczako K. (1996). Immunohistochemical evaluation of keratin 20 expression in intestinal metaplasia types I to III. *J. Clin. Pathol.* 49, 791-794.
- Simo P., Simon A.P., Bouziges F., Leberquier C., Keding M., Ekblom P. and Sorokin L. (1991). Changes in the expression of laminin during intestinal development. *Development* 112, 477-487.
- Simon-Assmann P., Duclos B., Orian-Rousseau V., Arnold C., Mathelin C., Engvall E. and Keding M. (1994). Differential expression of laminin isoforms in the developing human and mouse intestine. *Dev. Dyn.* 201, 71-85.
- Sipponen P., Kekki M. and Siurala M. (1991). The sydney system: epidemiology and natural history of chronic gastritis. *J. Gastroenterol. Hepatol.* 6, 244-251.
- Slomiany B.L., Piotrowski J., Sengupta S. and Slomiany A. (1991). Inhibition of gastric mucosal laminin receptor by *Helicobacter pylori* lipopolysaccharide. *Biochem. Biophys. Res. Commun.* 175, 77-81.
- Solcia E., Fiocca R., Luinetti O., Villani L., Padovan L., Calistri D., Ranzani G.N., Chiaravalli A. and Capella C. (1996). Intestinal and diffuse gastric cancers arise in a different background of *Helicobacter pylori* gastritis through different gene involvement. *Am. J. Surg. Pathol.* 20, S8-S22.
- Specian R.D. and Neutra M.R. (1984). Cytoskeleton of intestinal goblet cells in rabbit and monkey. *Theca. Gastroenterology* 87, 1313-1325.
- Spinard L., Einheber S., Cullen T., Milner T.A. and Giancotti F.G. (1995). Recombinant tail-less integrin beta 4 subunit disrupts hemidesmosomes but does not suppress alpha 6 beta 4 mediated cell adhesion to laminin. *J. Cell Biol.* 129, 473-487.
- Tani T., Karttunen T., Kiviluoto T., Kivilaakso E., Burgeson R.E., Sipponen P. and Virtanen I. (1996). $\alpha 6 \beta 4$ integrin and newly

- deposited laminin-1 and laminin-5 form the adhesion mechanism of gastric carcinoma. *Am. J. Pathol.* 149, 781-793.
- Tennenbaum T., Weiner A.K., Belanger A.J., Glick A.B., Hennings H. and Yuspa S.H. (1993). The suprabasal expression of $\alpha 6 \beta 4$ integrin is associated with a high risk for malignant progression in mouse skin carcinogenesis. *Cancer Res.* 53, 4803-4810.
- Timpl R., Rodhe H., Robey P.G., Rennard S.I., Foidart J.-M. and Martin G.R. (1979). Laminin-A glycoprotein from basement membranes. *J. Biol. Chem.* 254, 9933-9937.
- Tremblay E. and Ménard D. (1995). Compositional changes of laminin A chain isoforms, fibronectin and tenascin during the ontogenesis of human gastric mucosa. *Gastroenterology* 108, A758.
- Tremblay E. and Ménard D. (1996). Differential expression of extracellular matrix components during the morphogenesis of human gastric mucosa. *Anat. Rec.* 245, 668-676.
- Valkonen K.H., Wadström T. and Moran A.P. (1994). Interaction of lipopolysaccharides of *Helicobacter pylori* with basement membrane protein laminin. *Infect. Immun.* 62, 3640-3648.
- Virtanen I., Tani T., Bäck N., Häppölä O., Laitinen L., Kiviluoto T., Salo J., Burgeson R.E., Lehto V.P. and Kivilaakso E. (1995). Differential expression of laminin chains and their integrin receptors in human gastric mucosa. *Am. J. Pathol.* 147, 1123-1132.

Accepted May 25, 2001